

Parkinson's disease: Alpha synuclein, heme oxygenase and biotherapeutic countermeasures

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Bentham Science Publishers. Neurodegenerative disorders have been and remain persistent sources of enormous suffering throughout human history. The tragedy of their impact on human relationships, physical vitality, and fundamental dignity cannot be understated. Parkinson's disease (PD), one of the most common of these terrible illnesses, has a global incidence of approximately two-to-four percent of the human population, along with devastating social and economic impact. The present review analyzes aspects of PD pathophysiology that offer particularly attractive strategies for the development of improved prevention and therapy. The occurrence, symptoms, pathogenesis, and etiology of PD are considered, with focus on how the Alpha synuclein protein, which normally regulates neurotransmitter release, is aggregated by oxidative stressors into toxic inclusions, prominently including Lewy bodies and insoluble fibrils that disrupt the organization of brain areas responsible for motor control. The contribution to a progressively prooxidant tissue environment resulting from interaction between advanced glycation end products (AGEs) and their cognate receptors (RAGEs) is examined here as a significant driver of PD. This review also explores strategies currently being developed by a U.S.-Russian team that may reduce the risk and severity of PD by use of recombinant atoxic derivatives (ad) of botulinum neurotoxins (BoNT/A ad), that traffic inducers of the cytoprotective enzyme heme oxygenase to selected midbrain neurons, at which Alpha synuclein aggregation occurs. Considered together, the topic material presented here provides both researchers and clinicians with a short but concise overview of the current understanding of PD pathology and approaches to biotherapeutic (precision) countermeasures to its onset and progression.

<http://dx.doi.org/10.2174/1381612824666180717161338>

Keywords

Aggregation, Alpha synuclein, Heme oxygenase, Neurodegenerative disorders, Parkinson's disease, Proteotoxic stress

References

- [1] NIND. Parkinson's Disease Information Page 2017 [cited 2018 May 04.]. Available from: <https://www.ninds.nih.gov/disorders/All-Disorders/Parkinsons-Disease-Information-Page>.
- [2] Kalia LV, Lang AE. Parkinson's disease. *Lancet* 2015; 386(9996): 896-912.
- [3] Ling H, Massey LA, Lees AJ, Brown P, Day BL. Hypokinesia without decrement distinguishes progressive supranuclear palsy from Parkinson's disease. *Brain* 2012; 135(Pt 4): 1141-53.
- [4] Erro R, Stamelou M. The motor syndrome of Parkinson's disease. *Int Rev Neurobiol* 2017; 132: 25-32.
- [5] Samii A, Nutt JG, Ransom BR. Parkinson's disease. *Lancet* 2004; 363(9423): 1783-93.

- [6] Barranco Quintana JL, Allam MF, Del Castillo AS, Navajas RF. Parkinson's disease and tea: A quantitative review. *J Am Coll Nutr* 2009; 28(1): 1-6.
- [7] Alves RS, Barbosa ER, Scaff M. Postvaccinal parkinsonism. *Mov Disord* 1992; 7(2): 178-80.
- [8] Liu G, Zhang C, Yin J, Li X, Cheng F, Li Y, et al. alpha-Synuclein is differentially expressed in mitochondria from different rat brain regions and dose-dependently down-regulates complex I activity. *Neurosci Lett* 2009; 454(3): 187-92.
- [9] Sato K, Yamashita T, Kurata T, et al. Telmisartan reduces progressive oxidative stress and phosphorylated alpha-synuclein accumulation in stroke-resistant spontaneously hypertensive rats after transient middle cerebral artery occlusion. *J Stroke Cerebrovasc Dis* 2014; 23(6): 1554-63.
- [10] Lees AJ, Hardy J, Revesz T. Parkinson's disease. *Lancet* 2009; 373(9680): 2055-66.
- [11] Sveinbjornsdottir S. The clinical symptoms of Parkinson's disease. *J Neurochem* 2016; 139 (Suppl 1): 318-24.
- [12] Guo Q, You H, Yang X, Lin B, Zhu Z, Lu Z, et al. Functional single-walled carbon nanotubes 'CAR' for targeting dopamine delivery into the brain of Parkinsonian mice. *Nanoscale* 2017; 9(30): 10832-45.
- [13] Mahmoud FF, Al-Awadhi AM, Haines DD. Amelioration of human osteoarthritis symptoms with topical 'biotherapeutics': A phase I human trial. *Cell Stress Chaper* 2015; 20(2): 267-76.
- [14] Bitnun A, Ford-Jones EL, Petric M, MacGregor D, Heurter H, Nelson S, et al. Acute childhood encephalitis and *Mycoplasma pneumoniae*. *Clin Infect Dis* 2001; 32(12): 1674-84.
- [15] Chernov VM, Mouzykantov AA, Baranova NB, et al. Extracellular membrane vesicles secreted by mycoplasma *Acholeplasma laidlawii* PG8 are enriched in virulence proteins. *J Proteom* 2014; 110: 117-28.
- [16] Hu J, Chen C, Ou G, You X, Tan T, Hu X, et al. Nrf2 regulates the inflammatory response, including heme oxygenase-1 induction, by *Mycoplasma pneumoniae* lipid-associated membrane proteins in THP-1 cells. *Pathog Dis* 2017; 75(4).
- [17] Haines DD, Lekli I, Teissier P, Bak I, Tosaki A. Role of haeme oxygenase-1 in resolution of oxidative stress-related pathologies: focus on cardiovascular, lung, neurological and kidney disorders. *Acta Physiol* 2012; 204(4): 487-501.
- [18] Saxena S, Shukla D, Saxena S, et al. Hypoxia preconditioning by cobalt chloride enhances endurance performance and protects skeletal muscles from exercise-induced oxidative damage in rats. *Acta Physiol* 2010; 200(3): 249-63.
- [19] Csonka C, Varga E, Kovacs P, Ferdinandy P, Blasig IE, Szilvassy Z, et al. Heme oxygenase and cardiac function in ischemic/ reperfused rat hearts. *Free Radic Biol Med* 1999; 27(1-2): 119-26.
- [20] Balla J, Vercellotti GM, Jeney V, et al. Heme, heme oxygenase and ferritin in vascular endothelial cell injury. *Mol Nutr Food Res* 2005; 49(11): 1030-43.
- [21] Song W, Kothari V, Velly AM, et al. Evaluation of salivary heme oxygenase-1 as a potential biomarker of early Parkinson's disease. *Mov Disord* 2018; 33(4): 583-91.
- [22] Sun Y, Yang T, Leak RK, Chen J, Zhang F. Preventive and protective roles of dietary Nrf2 activators against central nervous system diseases. *CNS Neurol Disord Drug Targets* 2017; 16(3): 326-38.
- [23] Cuadrado A, Rojo AI. Heme oxygenase-1 as a therapeutic target in neurodegenerative diseases and brain infections. *Curr Pharm Des* 2008; 14(5): 429-42.
- [24] Taddei S, Bortolotto L. Unraveling the pivotal role of bradykinin in ACE inhibitor activity. *Am J Cardiovasc Drugs* 2016; 16(5): 309-21.
- [25] Lastres-Becker I, Ulusoy A, Innamorato NG, et al. alpha-Synuclein expression and Nrf2 deficiency cooperate to aggravate protein aggregation, neuronal death and inflammation in early-stage Parkinson's disease. *Hum Mol Genet* 2012; 21(14): 3173-92.
- [26] Nacereddine A, Bollacke A, Roka E, et al. Self-assembled supramolecular nanoparticles improve the cytotoxic efficacy of CK2 inhibitor THN7. *Pharmaceuticals* 2018; 11(1): 10.
- [27] Di Marco E, Gray SP, Jandeleit-Dahm K. Diabetes alters activation and repression of pro-and anti-inflammatory signaling pathways in the vasculature. *Front Endocrinol* 2013; 4: 68.
- [28] Vistoli G, De Maddis D, Cipak A, Zarkovic N, Carini M, Aldini G. Advanced glycoxidation and lipoxidation end products (AGEs and ALEs): an overview of their mechanisms of formation. *Free Radic Res* 2013; 47 (Suppl 1): 3-27.
- [29] Gao Y, Wake H, Morioka Y, et al. Phagocytosis of advanced glycation end products (AGEs) in macrophages induces cell apoptosis. *Oxidat Med Cell Long* 2017; 2017: 8419035.
- [30] Jiang X, Wang X, Tuo M, Ma J, Xie A. RAGE and its emerging role in the pathogenesis of Parkinson's disease. *Neurosci Lett* 2018; 672: 65-9.
- [31] Lin CH, Cheng YC, Nicol CJ, Lin KH, Yen CH, Chiang MC. Activation of AMPK is neuroprotective in the oxidative stress by advanced glycosylation end products in human neural stem cells. *Exp Cell Res* 2017; 359(2): 367-73.
- [32] Park EJ, Kim YM, Chang KC. Hemin reduces HMGB1 release by UVB in an AMPK/HO-1-dependent pathway in human keratinocytes HaCaT cells. *Arch Med Res* 2017; 48(5): 423-31.

- [33] Vazquez-Cintron EJ, Vakulenko M, Band PA, Stanker LH, Johnson EA, Ichtchenko K. Atoxic derivative of botulinum neurotoxin A as a prototype molecular vehicle for targeted delivery to the neuronal cytoplasm. *PLoS One* 2014; 9(1): e85517.
- [34] Vazquez-Cintron EJ, Beske PH, Tenezaca L, et al. Engineering botulinum neurotoxin C1 as a molecular vehicle for intra-neuronal drug delivery. *Sci Rep* 2017; 7: 42923.
- [35] Vazquez-Cintron E, Tenezaca L, Angeles C, et al. Pre-clinical study of a novel recombinant botulinum neurotoxin derivative engineered for improved safety. *Sci Rep* 2016; 6: 30429.